


Crop Report

Report name: [Block 510C] Crop report
 Report date: 12/07/2011
 Last climate date available: 10/07/2011
 Client name: EH Graham Centre
 Paddock name: Block 510C
 Report generated by: EH Graham Centre
 Date sown: 14-May
 Crop type: Wheat
 Variety sown: Wedgetail
 Sowing density: 89 plants/m²


Weather station used: Wagga Wagga AMO
 Rainfall records used: Weather station
 Soil type: Dirnaseer Red Kandosol
 Maximum rooting depth: 180 cm
 Roots constrained by EC:
 Stubble type: canola
 Stubble amount: 2500 kg/ha
 Start of growing season: 01-Apr
 Initial conditions date: 12-May
 Growing season rainfall to date: 89.6 mm
 Date of last rainfall entry: ?
 Expected harvest date: 26-Nov

Grain Yield Outcome



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture; the weather conditions so far; soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each of the past 100 years. The yield results are used to produce this graph.


Hay Yield Outcome



This graph show the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 335 kg/ha

The season so far - Growing Season Rainfall Deciles



GS11
emergence

GS12
2nd leaf

GS13
3rd leaf
1st tiller

GS14
4th leaf
early tillering

GS15
5th leaf
mid tillering

GS16
6th leaf
late tillering



Predicted

Earliest	24-May	5-Jun	17-Jun	28-Jun	7-Jul	17-Jul
Median	24-May	5-Jun	17-Jun	28-Jun	7-Jul	20-Jul
Latest	24-May	5-Jun	17-Jun	28-Jun	7-Jul	24-Jul



GS30
end of
tillering



GS31
1st node



GS32
2nd node



GS37
flag leaf



GS39
flag leaf
fully emerged



GS45
mid booting



GS55
mid head
emergence



GS65
mid
flowering



GS75
mid dough
fill

Predicted

Earliest	4-Sep	7-Sep	10-Sep	16-Sep	19-Sep	23-Sep	30-Sep	7-Oct	23-Oct
Median	10-Sep	13-Sep	16-Sep	24-Sep	27-Sep	2-Oct	8-Oct	15-Oct	31-Oct
Latest	16-Sep	19-Sep	22-Sep	3-Oct	6-Oct	11-Oct	18-Oct	26-Oct	15-Nov

Percentage of years in which frost occurs during flowering

Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	34%
Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	2%
Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	1%

Percentage of years in which heat shock occurs during grain fill (Z70-79)

Mild	
Maximum temperature between 32 and 34°C	43%
Moderate	
Maximum temperature between 34 and 36°C	25%
Severe	
Maximum temperature above 36°	10%


Incidence of frost for this growing season, during flowering

Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	0
Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0
Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0

Incidence of heat shock for this growing season, during grain fill (Z70-79)

Mild	
Maximum temperature between 32 and 34°C	0
Moderate	
Maximum temperature between 34 and 36°C	0
Severe	
Maximum temperature above 36°	0

Current distribution of PAW



PAW = Plant Available Water
CLL = Crop Lower Limit or Wilting Point
DUL = Drained Upper Limit or Field Capacity
PAWC = Plant Available Water Capacity
Current Crop PAW = Soil water currently accessible to the roots down to the current rooting depth
Soil PAW = Total accessible soil water in the soil profile

Water Budget

Initial PAW status @ 12-May	78 mm
Rainfall since 12-May	57 mm
Irrigations	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
Evaporation since 12-May	48 mm
Transpiration since 12-May	4 mm
Deep drainage since 12-May	0 mm
Run-off since 12-May	0 mm


Current PAW status: 84 mm

Nitrogen Budget


Initial N status @ 12-May	44 kg/ha
Mineralisation since 12-May	-9 kg/ha
N applications	14-May: 30 kg/ha
	8-Jul: 32 kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
Total N in plant	21 kg/ha
De-nitrification since 12-May	0 kg/ha
Leaching	0 kg/ha

Current N status: 76 kg/ha


Probability of Future Waterlogging Events




Current distribution of soil nitrogen (kg/ha)




Availability of Water to Growing Roots




Soil Nitrogen



Water Stress



Nitrogen Stress




Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Mean projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser.

Date	Growth Stage	Evap (mm)	Daily water use (mm)	Daily N use (kg/ha)	Water available to roots above stress threshold (mm)	Water available to roots above crop lower limit (mm)	N available to roots (kg/ha)
12-Jul	15.5	0.8	0.2	1.2	25.2	52.7	48.7
13-Jul	15.6	0.8	0.2	1.2	24.6	52.5	47.4
14-Jul	15.6	0.8	0.2	1.3	24.4	52.6	46.2
15-Jul	15.7	0.8	0.2	1.4	24.4	52.8	44.8
16-Jul	15.8	0.8	0.2	1.5	25.7	55.2	43.5
17-Jul	15.9	0.7	0.2	1.5	26.4	55.4	42.4
18-Jul	16.0	0.7	0.2	1.5	25.8	55.8	41.2
19-Jul	16.0	0.7	0.2	1.6	27.0	56.8	39.4
20-Jul	16.0	0.7	0.2	1.7	26.3	56.7	37.6
21-Jul	16.0	0.7	0.2	1.7	26.3	57.1	36.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

How much rainfall can I expect?
 The Bureau of Meteorology Forecast for the next 3 months




National Seasonal Rainfall Outlook: probabilities July to September 2011

Issued by the bureau of Meteorology 23rd June 2011

Queensland Department of Environment and Resource Management (DERM) 3 month rainfall forecast based on the current phase of the SOI

Probability of exceeding Median Rainfall



SOI Phase and analogue years


The SOI is currently in the Zero phase. The 30 day mean SOI for June was 0.9, in May it was 2.1.

The years in history with the same SOI phase:

1890, 1891, 1894, 1895, 1897, 1902, 1904, 1907, 1908, 1913, 1915, 1922, 1927, 1929, 1930, 1932, 1935, 1942, 1944, 1953, 1957, 1958, 1959, 1961, 1962, 1966, 1967, 1969, 1971, 1978, 1980, 1983, 1990, 1991, 2000, 2001, 2002, 2005, 2006, 2008, 2009

How much rainfall can I expect?


The SOI seasonal forecast for the next 3 months.



The SOI is an index that compares the atmospheric pressure between Tahiti and Darwin. SOI Phases are determined by comparing average monthly SOI values of the past two months. Phases of the SOI have been shown to be related to rainfall variability in a range of locations in Australia and around the world.

	Rainfall
Driest	0 to 101 mm
Drier	101 to 132 mm
Average	132 to 157 mm
Wetter	157 to 201 mm
Wettest	201 to 335 mm

How do seasonal forecasts affect yield expectations?



The 30 day mean SOI for June was 0.9, in May it was 2.1.

Yield outcomes of the current SOI Phase ARE NOT significantly different from yield outcomes of all years. Significance is determined on a 90% probability threshold. (PValue=0.651)