


Crop Report

Report name: Block 501W Crop report
 Report date: 22/07/2013
 Last climate date available: 20/07/2013
 Client name: EH Graham Centre
 Paddock name: Block 501W
 Report generated by: EH Graham Centre
 Date sown: 15-Apr
 Crop type: Wheat
 Variety sown: Wedgetail
 Sowing density: 70 plants/m²


Weather station used: Wagga Wagga
 Agricultural Institute
 Rainfall records used: Weather station
 Soil type: Red Kandosol (Dirnaseer No544)
 Maximum rooting depth: 100 cm
 Stubble type: Wheat
 Stubble amount: 4000 kg/ha
 Number of tillage operations: 0
 Stubble % incorporated into the top 10cm: 0 %
 Initial conditions date: 01-Apr
 Rainfall since 1-Apr: 159.5 mm
 Date of last rainfall entry: ?
 Expected maturity date: 17-Nov

Grain Yield Outcome



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture; the weather conditions so far; soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.


Hay Yield Outcome




This graph show the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 2616 kg/ha

The Season So Far - Growing Season Rainfall Deciles




Simulated and Predicted Crop Growth Stage



Predicted

	GS11 emergence	GS12 2nd leaf	GS13 3rd leaf 1st tiller	GS14 4th leaf early tillering	GS15 5th leaf mid tillering	GS16 6th leaf late tillering
Earliest	26-Apr	7-May	13-May	22-May	31-May	8-Jun
Median	26-Apr	7-May	13-May	22-May	31-May	8-Jun
Latest	26-Apr	7-May	13-May	22-May	31-May	8-Jun




Predicted

	GS30 end of tillering	GS31 1st node	GS32 2nd node	GS37 flag leaf	GS39 flag leaf fully emerged	GS45 mid booting	GS55 mid head emergence	GS65 mid flowering	GS75 mid dough fill
Earliest	21-Aug	24-Aug	28-Aug	2-Sep	5-Sep	10-Sep	18-Sep	23-Sep	11-Oct
Median	27-Aug	31-Aug	4-Sep	12-Sep	16-Sep	21-Sep	28-Sep	5-Oct	22-Oct
Latest	13-Sep	18-Sep	24-Sep	5-Oct	8-Oct	12-Oct	18-Oct	26-Oct	11-Nov

Probability and Incidence of Frost and Heat Shock

<i>Percentage of years in which frost occurs during flowering</i>		<i>Percentage of years in which heat shock occurs during grain fill (Z70-79)</i>	
Mild		Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	19%	Maximum temperature between 32 and 34°C	30%
Moderate		Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	4%	Maximum temperature between 34 and 36°C	13%
Severe		Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0%	Maximum temperature above 36°	7%
<i>Incidence of frost for this growing season, during flowering</i>		<i>Incidence of heat shock for this growing season, during grain fill (Z70-79)</i>	
Mild		Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	0	Maximum temperature between 32 and 34°C	0
Moderate		Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0	Maximum temperature between 34 and 36°C	0
Severe		Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0	Maximum temperature above 36°	0

Current Distribution of PAW



PAW = Plant Available Water
CLL = Crop Lower Limit or Wilting Point
DUL = Drained Upper Limit or Field Capacity
PAWC = Plant Available Water Capacity
Current Crop PAW = Soil water currently accessible to the roots down to the current rooting depth
Soil PAW = Total accessible soil water in the soil profile

Water Budget

Initial PAW status @ 1-Apr	43 mm
Rainfall since 1-Apr	159.5 mm
Irrigations	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
Evaporation since 1-Apr	45 mm
Transpiration since 1-Apr	29 mm
Deep drainage since 1-Apr	0 mm
Run-off since 1-Apr	3 mm

Current PAW status: 96 mm


Nitrogen Budget

Initial N status @ 01-Apr	200 kg/ha
N mineralisation since 01-Apr	7 kg/ha
N tie up since 01-Apr	13 kg/ha
N applications	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
Total N in plant	125 kg/ha
De-nitrification since 01-Apr	1 kg/ha
Leaching	0 kg/ha


Current N status: 68 kg/ha

Median N mineralisation to maturity = 6 kg/ha
 Median N tie up to maturity = 0 kg/ha


Probability of Future Waterlogging Events




Current distribution of soil nitrogen (kg/ha)




Availability of Water to Growing Roots




Availability of Soil Nitrogen to Growing Roots



Water Stress



Nitrogen Stress




Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Mean projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser.

Date	Growth Stage	Evap. (mm)	Water use (mm)	N use (kg/ha)	Water available to roots above stress threshold (mm)	Water available to roots above CLL (mm)	N available to roots (kg/ha)	Mineralisation (kg/ha)	N tie up (kg/ha)
22-Jul	16.0	0.2	0.6	1.3	52.0	95.0	41.6	0.1	0.0
23-Jul	16.0	0.2	0.8	1.3	51.7	94.7	40.3	0.0	0.0
24-Jul	16.0	0.2	0.7	1.2	52.4	95.4	39.1	0.0	0.0
25-Jul	16.0	0.2	0.8	1.1	52.2	95.2	37.9	0.0	0.0
26-Jul	16.0	0.2	0.7	1.1	51.3	94.4	36.8	0.0	0.0
27-Jul	16.0	0.2	0.7	1.0	51.8	94.8	35.8	0.0	0.0
28-Jul	16.0	0.2	0.7	1.0	52.6	95.6	34.8	0.0	0.0
29-Jul	16.0	0.2	0.8	0.9	52.1	95.2	33.8	0.0	0.0
30-Jul	16.0	0.2	0.8	0.9	51.5	94.5	32.9	0.0	0.0
31-Jul	16.0	0.2	0.7	0.8	52.7	95.7	32.1	0.0	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

How much rainfall can I expect?
 The Bureau of Meteorology Forecast for the next 3 months




National Seasonal Rainfall Outlook: probabilities July to September 2013

Issued by the bureau of Meteorology 19th June 2013

Queensland Department of Environment and Resource Management
 (DERM) 3 month rainfall forecast based on the current phase of the SOI

**Probability of exceeding
 Median Rainfall**



SOI Phase and analogue years


The SOI is currently in the Positive phase. The 30 day mean SOI for June was May was 10.6. In May the 31 day mean SOI was

The years in history with the same SOI phase:

1892, 1893, 1900, 1901, 1909, 1910, 1917, 1920, 1921, 1924, 1931, 1934, 1938, 1945, 1950, 1952, 1955, 1956, 1964, 1968, 1973, 1975, 1981, 1986, 1989, 1996, 1998

How much rainfall can I expect?


The SOI seasonal forecast for the next 3 months.



The SOI is an index that compares the atmospheric pressure between Tahiti and Darwin. SOI Phases are determined by comparing average monthly SOI values of the past two months. Phases of the SOI have been shown to be related to rainfall variability in a range of locations in Australia and around the world.

Rainfall	
Driest	0 to 100 mm
Drier	100 to 125 mm
Average	125 to 153 mm
Wetter	153 to 186 mm
Wettest	186 to 349 mm

How do seasonal forecasts affect yield expectations?



The 30 day mean SOI for June was May was 10.6. In May the 31 day mean SOI was 8.0.

Yield outcomes of the current SOI Phase ARE significantly different from yield outcomes of all years. Significance is determined on a 90% probability threshold. (PValue=0.029)