


# Crop Report

Report name: Crop Sequence High N Crop Report (Complete)  
 Report date: 21/07/2014  
 Last climate date available: 20/07/2014  
 Client name: EH Graham Centre  
 Paddock name: Crop Sequence High N  
 Report generated by: EH Graham Centre  
 Date sown: 15-May  
 Crop type: Wheat  
 Variety sown: Gregory  
 Sowing density: 150 plants/m<sup>2</sup>  
 Initial conditions date: 09-Apr  
 Soil type: Red Kandosol (No498-Generic)


SILO station used: Wagga Wagga AMO  
 Rainfall records used: SILO  
 Temperature records used: SILO  
 Maximum rooting depth: 100 cm  
 Stubble type: None  
 Stubble amount: kg/ha  
 Number of tillage operations: 0  
 Stubble % incorporated into the top 10cm: 0 %  
 Rainfall since 9-Apr: 185.2 mm  
 Date of last rainfall entry: ?  
 Expected maturity date: 23-Nov

## Grain Yield Outcome



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture; the weather conditions so far; soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.


## Hay Yield Outcome




This graph show the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 685 kg/ha

## The Season So Far - Growing Season Rainfall Deciles




## Simulated and Predicted Crop Growth Stage



### Predicted

	GS11 emergence	GS12 2nd leaf	GS13 3rd leaf 1st tiller	GS14 4th leaf early tillering	GS15 5th leaf mid tillering	GS16 6th leaf late tillering
<b>Earliest</b>	22-May	6-Jun	16-Jun	26-Jun	8-Jul	19-Jul
<b>Median</b>	22-May	6-Jun	16-Jun	26-Jun	8-Jul	19-Jul
<b>Latest</b>	22-May	6-Jun	16-Jun	26-Jun	8-Jul	19-Jul




### Predicted

	GS30 end of tillering	GS31 1st node	GS32 2nd node	GS37 flag leaf	GS39 flag leaf fully emerged	GS45 mid booting	GS55 mid head emergence	GS65 mid flowering	GS75 mid dough fill
<b>Earliest</b>	12-Aug	16-Aug	20-Aug	2-Sep	7-Sep	13-Sep	22-Sep	4-Oct	19-Oct
<b>Median</b>	18-Aug	22-Aug	26-Aug	8-Sep	13-Sep	21-Sep	2-Oct	11-Oct	28-Oct
<b>Latest</b>	22-Aug	26-Aug	31-Aug	14-Sep	19-Sep	30-Sep	11-Oct	21-Oct	11-Nov

## Probability and Incidence of Frost and Heat Shock

<i>Percentage of years in which frost occurs during flowering</i>		<i>Percentage of years in which heat shock occurs during grain fill (Z70-79)</i>	
<b>Mild</b>		<b>Mild</b>	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	40%	Maximum temperature between 32 and 34°C	39%
<b>Moderate</b>		<b>Moderate</b>	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	5%	Maximum temperature between 34 and 36°C	18%
<b>Severe</b>		<b>Severe</b>	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	1%	Maximum temperature above 36°	7%
<i>Incidence of frost for this growing season, during flowering</i>		<i>Incidence of heat shock for this growing season, during grain fill (Z70-79)</i>	
<b>Mild</b>		<b>Mild</b>	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	0	Maximum temperature between 32 and 34°C	0
<b>Moderate</b>		<b>Moderate</b>	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0	Maximum temperature between 34 and 36°C	0
<b>Severe</b>		<b>Severe</b>	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0	Maximum temperature above 36°	0

## Current Distribution of PAW



## Water Budget

Initial PAW status @ 9-Apr	3 mm
Rainfall since 9-Apr	185.2 mm
Irrigations	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
Evaporation since 9-Apr	81 mm
Transpiration since 9-Apr	6 mm
Deep drainage since 9-Apr	0 mm
Run-off since 9-Apr	11 mm

**Current PAW status: 83 mm**


## Nitrogen Budget

Initial N status @ 09-Apr	92 kg/ha
N mineralisation since 09-Apr	2 kg/ha
N tie up since 09-Apr	1 kg/ha
N applications	15-May: 25 kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
Total N in plant	37 kg/ha
De-nitrification since 09-Apr	3 kg/ha
Leaching	0 kg/ha


**Current N status: 80 kg/ha**

Median N mineralisation to maturity = 4 kg/ha  
 Median N tie up to maturity = 0 kg/ha


## Probability of Future Waterlogging Events




## Current distribution of soil nitrogen (kg/ha)




### Availability of Water to Growing Roots




### Availability of Soil Nitrogen to Growing Roots



### Water Stress



### Nitrogen Stress




Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

### Mean projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser.

Date	Growth Stage	Evap. (mm)	Water use (mm)	N use (kg/ha)	Water available to roots above stress threshold (mm)	Water available to roots above CLL (mm)	N available to roots (kg/ha)	Mineralisation (kg/ha)	N tie up (kg/ha)
21-Jul	16.0	0.7	0.3	1.6	54.9	79.9	64.5	0.0	0.0
22-Jul	16.0	0.7	0.3	1.8	54.3	79.4	63.0	0.0	0.0
23-Jul	16.0	0.8	0.4	1.9	55.2	80.4	61.1	0.0	0.0
24-Jul	16.0	0.7	0.4	1.9	55.7	81.0	59.4	0.0	0.0
25-Jul	16.0	0.7	0.4	2.2	56.0	81.2	57.2	0.0	0.0
26-Jul	16.0	0.7	0.4	2.1	55.6	81.0	55.2	0.0	0.0
27-Jul	16.0	0.6	0.4	2.2	56.4	81.7	53.1	0.0	0.0
28-Jul	16.0	0.6	0.5	2.3	57.7	83.2	50.8	0.0	0.0
29-Jul	16.0	0.6	0.5	2.3	56.9	82.4	48.5	0.0	0.0
30-Jul	16.0	0.6	0.5	2.3	57.0	82.6	46.4	0.0	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

How much rainfall can I expect?  
 The Bureau of Meteorology Forecast for the next 3 months




**National Seasonal Rainfall Outlook: probabilities July to September 2014**

Issued by the Bureau of Meteorology 25th June 2014.

For the outlook accuracy map go to: "[http://www.bom.gov.au/climate/ahead/rain\\_ahead.shtml#tabs=Outlook-accuracy](http://www.bom.gov.au/climate/ahead/rain_ahead.shtml#tabs=Outlook-accuracy)"

Queensland Department of Environment and Resource Management (DERM) 3 month rainfall forecast based on the current phase of the SOI

**Probability of exceeding Median Rainfall**



## SOI Phase and analogue years


The SOI is currently in the Zero phase. The 30 day mean SOI for June was -0.8. In May the 31 day mean was 4.3.

The years in history with the same SOI phase:

1890, 1891, 1894, 1895, 1897, 1902, 1904, 1907, 1908, 1913, 1915, 1922, 1927, 1929, 1930, 1932, 1935, 1942, 1944, 1953, 1957, 1958, 1959, 1961, 1962, 1966, 1967, 1969, 1971, 1978, 1980, 1983, 1990, 1991, 2000, 2001, 2002, 2005, 2006, 2008, 2009

## How much rainfall can I expect?


The SOI seasonal forecast for the next 3 months.



The SOI is an index that compares the atmospheric pressure between Tahiti and Darwin. SOI Phases are determined by comparing average monthly SOI values of the past two months. Phases of the SOI have been shown to be related to rainfall variability in a range of locations in Australia and around the world.

	Rainfall
Driest	0 to 97 mm
Drier	97 to 131 mm
Average	131 to 158 mm
Wetter	158 to 202 mm
Wettest	202 to 347 mm

## How do seasonal forecasts affect yield expectations?



The 30 day mean SOI for June was -0.8. In May the 31 day mean was 4.3.

Yield outcomes of the current SOI Phase ARE NOT significantly different from yield outcomes of all years. Significance is determined on a 90% probability threshold. (PValue=0.277)