


Crop Report

Report name: Crop Sequence High N Crop Report (Complete)
 Report date: 09/07/2014
 Last climate date available: 7/07/2014
 Client name: EH Graham Centre
 Paddock name: Crop Sequence High N
 Report generated by: EH Graham Centre
 Date sown: 15-May
 Crop type: Wheat
 Variety sown: Gregory
 Sowing density: 150 plants/m²
 Initial conditions date: 09-Apr
 Soil type: Red Kandosol (No498-Generic)


SILO station used: Wagga Wagga AMO
 Rainfall records used: SILO
 Temperature records used: SILO
 Maximum rooting depth: 100 cm
 Stubble type: None
 Stubble amount: kg/ha
 Number of tillage operations: 0
 Stubble % incorporated into the top 10cm: 0 %
 Rainfall since 9-Apr: 166.6 mm
 Date of last rainfall entry: ?
 Expected maturity date: 23-Nov

Grain Yield Outcome



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture; the weather conditions so far; soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.


Hay Yield Outcome




This graph show the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 366 kg/ha

The Season So Far - Growing Season Rainfall Deciles




Simulated and Predicted Crop Growth Stage



Predicted

	GS11 emergence	GS12 2nd leaf	GS13 3rd leaf 1st tiller	GS14 4th leaf early tillering	GS15 5th leaf mid tillering	GS16 6th leaf late tillering
Earliest	22-May	6-Jun	16-Jun	26-Jun	8-Jul	16-Jul
Median	22-May	6-Jun	16-Jun	26-Jun	8-Jul	21-Jul
Latest	22-May	6-Jun	16-Jun	26-Jun	8-Jul	25-Jul



Predicted

	GS30 end of tillering	GS31 1st node	GS32 2nd node	GS37 flag leaf	GS39 flag leaf fully emerged	GS45 mid booting	GS55 mid head emergence	GS65 mid flowering	GS75 mid dough fill
Earliest	12-Aug	16-Aug	21-Aug	2-Sep	5-Sep	12-Sep	23-Sep	2-Oct	19-Oct
Median	19-Aug	22-Aug	27-Aug	8-Sep	13-Sep	21-Sep	2-Oct	11-Oct	28-Oct
Latest	25-Aug	28-Aug	2-Sep	15-Sep	21-Sep	1-Oct	12-Oct	22-Oct	12-Nov

Probability and Incidence of Frost and Heat Shock

Percentage of years in which frost occurs during flowering

Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	38%
Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	5%
Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	1%

Percentage of years in which heat shock occurs during grain fill (Z70-79)

Mild	
Maximum temperature between 32 and 34°C	39%
Moderate	
Maximum temperature between 34 and 36°C	18%
Severe	
Maximum temperature above 36°	8%


Incidence of frost for this growing season, during flowering

Mild	
Minimum temperature between 2 and 0°C during flowering (Z60-69)	0
Moderate	
Minimum temperature between 0 and -2°C during flowering and early grain fill (Z60-75)	0
Severe	
Minimum temperature less than -2°C during flowering and grain fill (Z60-79)	0

Incidence of heat shock for this growing season, during grain fill (Z70-79)

Mild	
Maximum temperature between 32 and 34°C	0
Moderate	
Maximum temperature between 34 and 36°C	0
Severe	
Maximum temperature above 36°	0

Current Distribution of PAW



Water Budget

Initial PAW status @ 9-Apr	3 mm
Rainfall since 9-Apr	166.6 mm
Irrigations	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
	: mm
Evaporation since 9-Apr	71 mm
Transpiration since 9-Apr	3 mm
Deep drainage since 9-Apr	0 mm
Run-off since 9-Apr	11 mm

Current PAW status: 77 mm


Nitrogen Budget

Initial N status @ 09-Apr	92 kg/ha
N mineralisation since 09-Apr	2 kg/ha
N tie up since 09-Apr	1 kg/ha
N applications	15-May: 25 kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
	: kg/ha
Total N in plant	21 kg/ha
De-nitrification since 09-Apr	3 kg/ha
Leaching	0 kg/ha


Current N status: 96 kg/ha

Median N mineralisation to maturity = 4 kg/ha
 Median N tie up to maturity = 0 kg/ha

Probability of Future Waterlogging Events




Current distribution of soil nitrogen (kg/ha)




Current Crop Available N = 79 kg/ha
 Total Soil N = 96 kg/ha


Availability of Water to Growing Roots




Availability of Soil Nitrogen to Growing Roots



Water Stress



Nitrogen Stress




Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Mean projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser.

Date	Growth Stage	Evap. (mm)	Water use (mm)	N use (kg/ha)	Water available to roots above stress threshold (mm)	Water available to roots above CLL (mm)	N available to roots (kg/ha)	Mineralisation (kg/ha)	N tie up (kg/ha)
9-Jul	15.2	0.7	0.2	1.1	47.4	70.2	76.5	0.0	0.0
10-Jul	15.3	0.7	0.2	1.1	46.9	69.9	75.6	0.0	0.0
11-Jul	15.3	0.6	0.2	1.2	47.7	70.8	74.7	0.0	0.0
12-Jul	15.4	0.7	0.2	1.2	47.2	70.5	74.0	0.0	0.0
13-Jul	15.5	0.7	0.2	1.2	47.3	70.9	73.1	0.0	0.0
14-Jul	15.6	0.7	0.2	1.3	47.8	71.8	71.9	0.0	0.0
15-Jul	15.7	0.7	0.2	1.4	50.2	74.2	70.9	0.0	0.0
16-Jul	15.8	0.7	0.2	1.5	52.1	76.3	69.7	0.0	0.0
17-Jul	15.8	0.6	0.2	1.5	53.0	77.4	68.7	0.0	0.0
18-Jul	15.9	0.7	0.2	1.5	53.2	77.5	67.4	0.0	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

How much rainfall can I expect?
 The Bureau of Meteorology Forecast for the next 3 months




National Seasonal Rainfall Outlook: probabilities July to September 2014

Issued by the Bureau of Meteorology 25th June 2014.

For the outlook accuracy map go to: "http://www.bom.gov.au/climate/ahead/rain_ahead.shtml#tabs=Outlook-accuracy"

Queensland Department of Environment and Resource Management (DERM) 3 month rainfall forecast based on the current phase of the SOI

Probability of exceeding Median Rainfall



July / September based on consistently near zero phase during May / June



SOI Phase and analogue years


The SOI is currently in the Zero phase. The 30 day mean SOI for June was -0.8. In May the 31 day mean was 4.3.

The years in history with the same SOI phase:

1890, 1891, 1894, 1895, 1897, 1902, 1904, 1907, 1908, 1913, 1915, 1922, 1927, 1929, 1930, 1932, 1935, 1942, 1944, 1953, 1957, 1958, 1959, 1961, 1962, 1966, 1967, 1969, 1971, 1978, 1980, 1983, 1990, 1991, 2000, 2001, 2002, 2005, 2006, 2008, 2009

How much rainfall can I expect?


The SOI seasonal forecast for the next 3 months.



The SOI is an index that compares the atmospheric pressure between Tahiti and Darwin. SOI Phases are determined by comparing average monthly SOI values of the past two months. Phases of the SOI have been shown to be related to rainfall variability in a range of locations in Australia and around the world.

	Rainfall
Driest	0 to 97 mm
Drier	97 to 131 mm
Average	131 to 158 mm
Wetter	158 to 202 mm
Wettest	202 to 347 mm

How do seasonal forecasts affect yield expectations?



The 30 day mean SOI for June was -0.8. In May the 31 day mean was 4.3.

Yield outcomes of the current SOI Phase ARE NOT significantly different from yield outcomes of all years. Significance is determined on a 90% probability threshold. (PValue=0.276)